


# Adult Cardiac Arrest Circular Algorithm



## CPR Quality

- Push hard (at least 2 inches [5 cm]) and fast (100-120/min) and allow complete chest recoil.
- Minimize interruptions in compressions.
- Avoid excessive ventilation.
- Change compressor every 2 minutes, or sooner if fatigued.
- If no advanced airway, 30:2 compression-ventilation ratio.
- Quantitative waveform capnography
  - If PETCO<sub>2</sub> is low or decreasing, reassess CPR quality.

## Shock Energy for Defibrillation

- **Biphasic:** Manufacturer recommendation (eg, initial dose of 120-200 J); if unknown, use maximum available. Second and subsequent doses should be equivalent, and higher doses may be considered.
- **Monophasic:** 360 J

## Drug Therapy

- **Epinephrine IV/IO dose:** 1 mg every 3-5 minutes
- **Amiodarone IV/IO dose:** First dose: 300 mg bolus. Second dose: 150 mg.
- *or*
- **Lidocaine IV/IO dose:** First dose: 1-1.5 mg/kg. Second dose: 0.5-0.75 mg/kg.

## Advanced Airway

- Endotracheal intubation or supraglottic advanced airway
- Waveform capnography or capnometry to confirm and monitor ET tube placement
- Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions

## Return of Spontaneous Circulation (ROSC)

- Pulse and blood pressure
- Abrupt sustained increase in PETCO<sub>2</sub> (typically ≥40 mm Hg)
- Spontaneous arterial pressure waves with intra-arterial monitoring

## Reversible Causes

- |                           |                         |
|---------------------------|-------------------------|
| • Hypovolemia             | • Tension pneumothorax  |
| • Hypoxia                 | • Tamponade, cardiac    |
| • Hydrogen ion (acidosis) | • Toxins                |
| • Hypo-/hyperkalemia      | • Thrombosis, pulmonary |
| • Hypothermia             | • Thrombosis, coronary  |