

# Change Notice

## *ACLS Provider Manual*

Print ISBN 978-1-68472-295-2, AHA Product Number 25-1106

eBook ISBN 978-1-68472-306-5, AHA Product Number 25-3100

| Print Page Number | Location                                                                                        | Original Text                                                                                                                                                                                                                        | Change                                                                                                                                                                        | When Change Was Identified |
|-------------------|-------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------|
| 148               | Figure 60. Adult and Pediatric Durable LVAD Algorithm > Assessing Perfusion table > Last bullet | PETCO <sub>2</sub> >20 mm Hg (if available and should be used only when an ET tube or tracheostomy is used to ventilate the patient; use of a supraglottic [eg, King] airway results in a falsely elevated PETCO <sub>2</sub> value) | PETCO <sub>2</sub> >20 mm Hg                                                                                                                                                  | 10/22/25                   |
| 128               | Adult Cardiac Arrest Circular Algorithm > table > Advanced Airway cell > Delete 3rd bullet      | <ul style="list-style-type: none"> <li>Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions</li> </ul>                                                                   | <ul style="list-style-type: none"> <li><del>Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions</del></li> </ul> | 10/29/25                   |
| 189               | Science Summary Table > ACLS topic column, ACS row > 2 <sup>nd</sup> bullet                     | <ul style="list-style-type: none"> <li>Removing clopidogrel as primary anticoagulant</li> </ul>                                                                                                                                      | <ul style="list-style-type: none"> <li>Removing clopidogrel as primary antiplatelet</li> </ul>                                                                                | 10/29/25                   |