

Change Notice

ACLS Provider Manual

Print ISBN 978-1-68472-295-2, AHA Product Number 25-1106
 eBook ISBN 978-1-68472-306-5, AHA Product Number 25-3100

Print Page Number	Location	Original Text	Change	When Change Was Identified
148	Figure 60. Adult and Pediatric Durable LVAD Algorithm > Assessing Perfusion table > Last bullet	PETCO ₂ >20 mm Hg (if available and should be used only when an ET tube or tracheostomy is used to ventilate the patient; use of a supraglottic [eg, King] airway results in a falsely elevated PETCO ₂ value)	PETCO ₂ >20 mm Hg	10/22/25
128	Adult Cardiac Arrest Circular Algorithm > table > Advanced Airway cell > Delete 3rd bullet	• Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions	• Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions	10/29/25
189	Science Summary Table > ACLS topic column, ACS row > 2 nd bullet	• Removing clopidogrel as primary anticoagulant	• Removing clopidogrel as primary antiplatelet	10/29/25
105	Part 3: High-Performance Teams > Respiratory Arrest > Identifying Respiratory Problems by Severity > Respiratory Arrest > 2 nd paragraph > 3 rd sentence, last word > change “risk” to “rise”	“...delivered to produce chest risk.”	“...delivered to produce chest rise.”	01/26/26
164	Part 3: High-Performance Teams > Post-Cardiac Arrest Care > Neuroprognostication > Figure 67. Schematic overview of a multimodal approach to neuroprognostication > Abbreviation list > Change “electrocardiogram” to “electroencephalogram”	“...EEG, electrocardiogram; ...”	“...EEG, electroencephalogram; ...”	01/26/26