

Change Notice

Heartsaver Pediatric First Aid CPR AED Student Workbook

Print ISBN 978-1-61669-425-8, AHA Product Number 15-1038

eBook ISBN 978-1-61669-510-1, AHA Product Number 15-3113

Print Page Number	Location	Original Text	Change	When Change Was Made
38	Part 2: under “Use an Epinephrine Pen,” in the table “How to Use an Epinephrine Pen,” in the third row	Press the tip of the injector...	Hold the leg firmly in place just before and during the injection. Press the tip of the injector...	At next printing or update after 4/28/2017
38	Part 2: under “Use an Epinephrine Pen,” in the table “How to Use an Epinephrine Pen,” in the fourth row	Hold the pen in place for about 10 seconds.	For EpiPen and EpiPen Jr injectors, hold the injector in place for 3 seconds. Some other injectors may be held in place for up to 10 seconds. Be familiar with the manufacturer’s instructions for the type of injector you are using.	At next printing or update after 4/28/2017
44	Part 2, under “Diabetes and Low Blood Sugar,” the table in the section “Step 4: Act”		{Replace the table with the table at the end of this document.}	At next printing or update after 10/24/2016
159	CPR AED, under “CPR for Infants,” in the section “Definition of an Infant”	from 1 month to 1 year of age	less than 1 year old	At next printing or update after 10/24/2016
183	In the green row at the top of the table, under the word “Infants”	(Age Less Than 1 Year, Excluding Newborns)	(Age Less Than 1 Year)	At next printing or update after 10/24/2016
184	In the green row at the top of the table, under the word “Infants”	(Age Less Than 1 Year, Excluding Newborns)	(Age Less Than 1 Year)	At next printing or update after 10/24/2016

How to Help a Child With Low Blood Sugar

Check the child's First Aid Action Plan. Follow the plan, including directions about how to check blood sugar.

If the child can't sit up and swallow

Phone or have someone phone 9-1-1. Do not try to give the child anything to eat or drink.

If the child can sit up and swallow

Give the child something that contains sugar to eat or drink.

Have the child sit quietly or lie down.

If the child does not improve within 15 minutes, phone or have someone phone 9-1-1.